



## ***Diagnosis of The Soil-Transmitted Parasites in Wasit Province-Iraq***

**Noor Naeem Shakir, Suadad B. Khairi**

*Department of Biology, College of Education for Pure Science, Wasit university, IRAQ*

### **Article Information**

Received: 11-2-2026

Accepted: 1-3-2026

Published: 1-4-2026

### **Abstract**

Such places of leisure as the dirt found in parks and public places can be infected with the stray animals in these gardens, which can lead to contagious stages of parasites. A great number of individuals sleep in them and could be exposed to infection by the parasites since the parasites' infectious phases are able to endure years or even months. regardless of the weather. This is the reason why the study was carried out in various distinct locations of Wasit Governorate.

Simple random selection was used to collect soil samples which were used to collect a total of 35 soil samples at different locations in Wasit and the modified flotation technique was used to process them. The overall parasite contamination percentage of all the studied soil samples was 86 ( 30 / 35). Eight different forms of helminthes were identified from all of the soil samples that were examined. (35 ) *Toxocara* and *Ascaris ova* had the highest frequently occurring helminthes in 28% (20 ) of the samples followed by of *Ancylostoma* in 14 percent (10 ), *Trichuris* in 3 percent ( 2 ),*Enterobius* in 6 percent (4 ), *Fasciola* metacercariae in 3 percent (2 ),*Diphyllobothrium* in 6 percent ( 4 ).The results indicate that the communities that are at risk of zoonotic transmission are those that are at parks and other places of community.

**Keywords:** Soil transmitted helminths: STH: soil contamination: Flotation technique

### **Introduction**

Helminths also known as soil-transmitted helminths or STHs, are a diverse group of intestinal parasites. the causative agents of some of the most prevalent intestinal infections, in particular, in the world [Caldrer *et al.*, 2020; Loukas *et al.*, 2021]. The most widespread ones are *Ascaris lumbricoides*, *Trichloroe* thene *Trichuris trichiura*, hookworms, including *Ancylostoma duodenale* and *Necator americanus* [Nath *et al.*, 2020 ; Raj *et al.*, 2022]. These parasites are similar in terms of methods of transmission: their eggs or larvae are present in the feces of sick people and pollute the environment, and in particular, the soil. These

eggs can take very long durations of time in warm and damp conditions hence a source of constant infection. STH infections can cause many health problems, including undernourishment, anemia, stunted growth, and delayed cognitive development in kids [Caldrer *et al.*, 2020; Masangcay *et al.*, 2021; Scott and Koski, 2021].

This health impact and economic cost associated with the parasitic helminths, i.e. Human and animal soil-transmitted helminths (STHs) constitutes a significant burden of illnesses in terms of morbidity and mortality, particularly in the disadvantaged populations with minimal resources, which has a significant impact on The worldwide chain of food handling. The World Health Organization (WHO) states that the most prevalent STH species in humans are *Ascaris lumbricoides*, *Trichloro trichiura*, and hookworms *Ancylostoma duodenale* and *Necator americanus*. [WHO, 2023].

The poor environmental conditions, absence of sources of clean water, as well as low socioeconomic standing, which are the characteristic features of The high prevalence indexes in low- and middle-income nations are predisposed by their populations [Ihnacik *et al.*, 2022; Montresor *et al.*, 2020]. In addition to that, these zoonotic agents are distributed depending on the Soil types, wet and warm soils and the local conditions [Afonso *et al.*, 2008; Collender *et al.*, 2015; Thevenet *et al.*, 2020; Umhang *et al.*, 2017]. The stages of the parasite are also encouraged to develop throughout tropical and subtropical regions of the world and directly infect diseased people or animals [Ihnacik *et al.*, 2022]. Geophagia and placing dirty hands in the mouth also become risk factors [Otero *et al.*, 2018; Otranto and Deplazes, 2019; Traversa *et al.*, 2014]. Therefore, the congestion of the beach, park, backyard or sandbox with zoonotic parasite poses a health risk to people [Traversa *et al.*, 2014; Eisen *et al.*, 2019]. There is also a growing realization that zoonotic diseases can be transmitted by eating of raw and fresh foodstuff, which has been exposed to the foodborne parasites contaminated soil [Koutsoumanis *et al.*, 2018; Oranto and Deplazes, 2019; Utaaker *et al.*, 2017].

Even though these parasitic infections don't kill many people, They can result in several health issues and impairments [Gosha *et al.*, 2012] in vulnerable populations such as women of reproductive age (WRA), school-aged children (SAC), and preschool-aged children (PSAC) [Zeynudin *et al.*, 2022] and young animals [Kalkal *et al.*, 2020]. This means that a lot of work needs to be done to control them. The condition is thought to be responsible for 3.4 million disability-adjusted life years (DALYs) globally, with at least one of these STH species believed to infect over one-fifth of the population [Jourdan *et al.*, 2018; WHO, 2018].

The current study aims to ascertain the Wasit province's soil-transmitted helminthes (STH) infection prevalence.

## Methods

### 1. Soil collection and processing

In the year 2024- 2025; 35 soil samples were picked randomly in different locations in Wasit. Initially, the town was physically divided into five areas, which were north, south, east, west and center. Each region had thirty samples. The research was based on parks, communal areas. About 50 g were taken out of 3 cm depth of the ground per collection. Since some of the samples were moist, hence all the samples were dried at room temperature and in an air-dry in approximately 24 h on tray.

## 2. Sucrose flotation method

A modified flotation technique was used to process soil samples [Horiuchi and Uga ,2016]. In short, a 150-um sieve was used to sift the soil samples after they had been air dried for 24 to 48 hours. A vortex mixer was used to combine three grams of sieved soil with eight milliliters of distilled water in a ten milliliter test tube. After that, the suspension was centrifuged for five minutes at 1000 rpm. After removing the supernatant, we filled the tube with 8 milliliters of sucrose solution with a specific gravity of 1.2. Then we stirred it well with a vortex mixer. The tube was centrifuged for  $800 \times g$  for 5 min. After centrifugation, sucrose solution at 1.3 specific gravity was slowly added up to the brim of the tube using a 10-ml syringe until an upper meniscus was formed. To gather the uppermost part of the sucrose suspension, a cover slip was carefully positioned on the meniscus. STH eggs were detected on the slides using a compound light microscope set to  $100\times$  and  $400\times$  magnification

### Statistical Analysis

The Statistical Package for the Social Sciences (SPSS), version 25.0 (IMB Corp, Armonk, New York, USA), was used to analyze the data. Achi-squared test was used to compare the associations between proportions .  $P < 0.050$  was considered statistically significant .


## Results

The percent contamination of all the soil samples analyzed with parasites was 86% (30 / 35). As per the types of helminthes 8 types were reported, out of the total number of sampled soils (35 ), *Toxocara* and *Ascaris* ova were the most common helminthes in 20 samples 28% , followed by *Ancylostoma* with 10 positive 14% , *Trichuris* 2 (3%),*Enterobius* 4 (6%), *Fasciola* metacercariae 2( 3%), *Diphyllobothrium* 4(6%) , *Strongyloides* 10(14%). as reported in Table (1 ) and figure(1)

**Table (1 ): Total numbers and percentages of helminthic genera' eggs contaminating soil (n = 35)**

Type of helminthes	No. of contaminated soil	samples Percentage
<i>Toxocara</i>	20	28%
<i>Ascaris</i>	20	28%
<i>Ancylostoma</i>	10	14%
<i>Trichuris</i>	2	3%
<i>Enterobius</i>	4	6%
<i>F. metacercariae</i>	2	3%
<i>Diphyllobothrium</i>	4	6%
<i>Strongyloides</i>	10	14%
<i>P value</i>	0.001**	


Chi-square test; \*\*: significant at  $P < 0.05$ .



**Fig (1): Typical pictures of the parasites discovered during this investigation, 1: hook egg 2: *Toxocara*, 3 *Trichuris* 4: *Ascaris lumbricoides* 5: *Strongyloides* 7: *Enterobius***

The overall parasite contamination level of all soil samples analyzed was 86% ( 30 / 35). The parasite contamination rate of the soil samples, which studied in Kut, was the highest, at 19% and the lowest contamination rate was 6% in the Al Numaniya district. The occurrence of parasites in the soils of Badra ,Al Hayy, Sheak saad, Al Dujaly and Al Muwafaqia were 11% 14% 13% 17% and 21%, respectively.

In Kut, eight different species of helminthic ova were identified from contaminated soil samples taken from several areas in the Wasit Governorate (Table 4), with *Ascaris* having the highest rate at 14% of the samples, followed by *Ancylostoma* 3(21%), *Strongyloides* 4(29%), *Trichuris* 1(7%), *Toxocara* 1(7%), *Enterobius* 1(7%), *Diphyllobothrium* 1(7%) , *Fasciola metacercariae* 1(7%).



**Figure (2) : Pie chart showing the frequency distribution of soil pollution by helminthic genus' eggs.**

The Badra district's soil samples were analyzed, and the findings showed that *Ascaris* and *Toxocara* had the highest percentage of contamination (38%), followed by *Strongyloides* 1 (13%) *Toxocara* 3(38%) and *Fasciola metacercariae* 1 (13%). The results showed that *Toxocara* had the highest rate of contamination in the Al-Numaniya district at 75% (3), followed by *Ascaris* 1 at 25%. In Al- Hayy district, the findings showed that the highest pollution rate was with *Toxocara* at 40 % (4), followed by *Ascaris* 30% (3), followed by *Ancylostoma*, *Strongyloides* and *Enterobius* the same rate of 10% for each of them. In Sheak saad district, the findings showed that the highest pollution rate was with *Ascaris* 4(44%) followed by *Toxocara* 22% and *Ancylostoma*, *Enterobius* *Diphyllobothrium* the same rate of 11% for each of them, and in Al- dujaly the results revealed that the highest rate of contamination was with *Ascaris* and *toxocara* 4(33%) followed by *Ancylostoma* 2(17%), *Enterobius* *Diphyllobothrium* the same rate of 8% (1) for each, In Al- muwafaqia The findings showed that the greatest contamination rate was with *Strongyloides* 4(27%) followed by *Ascaris*, *toxocara* and *Ancylostoma* 3(20%), *Diphyllobothrium* and *Trichuris* the same rate of 1 (7%) for each. table (2).

**Table (2): Soils contamination of Helminthes type versus Districts -Wasit province.**

Districts							
Contaminated with parasite types	Kut	Badra	Al-Numaniya	Al-Hayy	Sheak saad	Al-dujaly	Al-muwafaqia
<i>Ascaris</i>	29 (14.3%)	3(37.5%)	1(25%)	3(30%)	4(44.4%)	4 (33.3%)	3 (20%)
<i>Ancylostoma</i>	3 (21.4%)	0	0	1(10%)	1(11.1%)	2 (16.7%)	3 (20%)
<i>Strongyloides</i>	4 (28.6%)	1 (12.5%)	0	1(10%)	0	0	4 (26.7%)
<i>Trichuris</i>	1 (7.1%)	0	0	0	0	0	1 (6.7%)
<i>Toxocara</i>	1 (7.1%)	3(37.5)	3(75%)	4(40%)	2 (22.2%)	4 (33.3%)	3 (20%)
<i>Enterobius</i>	1 (7.1%)	0	0	1(10%)	1 (11.1%)	1 (8.3%)	0
<i>Diphyllobothrium</i>	1 (7.1%)	0	0	0	1 (11.1%)	1 (8.3%)	1 (6.7%)
<i>F. metacercariae</i>	1 (7.1%)	1(12.5%)	0	0	0	0	0
<b>Total</b>	14	8	4	10	9	12	15
<b>%</b>	19%	11%	6%	14%	13%	17%	21%
<b>P value</b>	0.608	0.572	0.180	0.406	0.437	0.429	0.700

Chi-square test; \*\*: significant at  $P < 0.05$ .

## Discussion

The presence of soil contamination with parasite is a positive indication of the threat of transmission of the parasites to human beings or animals. In the present research, the total contamination of soil specimens was 86 percentage of 30 specimens (30/35) of analyzed specimens and this rate is a predictive indicator of a potential threat to the public health due to such ova. FKUI (2016) asserts that the habitat of Soil transmitted helminths worm eggs is to infect the sections of the small intestine. This parasite is cosmopolitan in its occurrence particularly in the humid air of the area and is highly associated to sanitary and hygiene conditions.

The result is similar to those of other researchers across the globe which discovered higher concentrations of helminthic and helminthic ova in soil. Rates of 67%, 68.1%; 50.6 and 35% were reported in Sudan, Nigeria, and Ethiopia respectively [Mohammed *et al.*, 2016 ; Hasan *et al.*, 2017 ; Badaki *et al.*, 2018; Tadege *et al.*, 2022]. In comparison, south Cameroon had a reduced rate of 17.6% STH [Golek and Al-Saeed , 2019 ].

Our study also agrees with studies conducted inside Iraq such as: In Duhok Governorate, Iraq, 593 (84.71%) of the 700 soil samples that were gathered from various locations were detected in the soil samples that were evaluated. The most common helminths were *Taenia* eggs, which were found in 117 (16.71%), followed by *Hymenolepis nana* ova in 107 (15.28%), *Toxocara* ova in 106 (15.14%), *Ancylostoma* ova in 95 (13.57%), *Ascaris* ova in 71 (10.14%), *Hymenolepis diminuta* ova in 51 (7.28%) and *Trichuris* ova in 46 (6.57%) [Khairi *et al.*, 2022 ]. These results agree with the results of [34] It is present in veggies gathered from various marketplaces in Wasit province /Iraq Out of 300 samples of the six types of vegetables , 247 were contaminated by parasite with rate (82.33%), *Strongyloides stercoralis* was the most often found parasite with (100) , *Ascaris lumbricoides* (85) , *Iodamoeba butschlii* (76) and *Ancylostoma duodenale*(59), *Enterobius vermicularis* (50), *Fasciola hepatica* (19) and *Hookworm* egg (17) , A lower rate of infection was reported for *Hymenolepis nana*, *Balantidium coli*, and *Diphyllobothrium latum*, with 8, 7, and 2 cases, respectively.

The environment, closeness to latrines, availability of trash, and high vegetation cover of the sampling areas were some of the environmental factors that played a significant role in the survival of STH eggs in the Wasit soil. These findings support the results of [Tadege *et al.*, 2022 ] who emphasized that humid and shady soils promote the survival of helminth eggs. FKUI (2016) also highlighted the fact that STH eggs can take months under humid soils that are not exposed to ultraviolet sunlight that decays the viability of eggs. The warm climate with high humidity in Sindupraja provides the best environment to maintain eggs and become infected, and locations with higher sanitation and sunlight exposure, seen by [Anwar *et al.*, 2020] are less contaminated with soil.

These disparities are mainly affected by the differences in sanitation, population density and hygiene practices. This is particularly common in and around the riverbanks and other areas with poor sanitation, which further confirms the interaction between the soil moisture and poor hygiene and the presence of parasites.

## Conclusion

The findings of our study in conclusions show that there is a high contamination of soil by the egg and helminths parasites in the Wasit province and particularly in the parks and the areas of public places. Moreover, it was discovered that these helminthic can infect human beings and animals which in turn can

lead to diseases and bring numerous health complications to them. This survey is the first survey on *Toxocara*, *Ascaris*, *Ancylostoma*, *Trichuris*, *Enterobius*, *Fasciola metacercariae*, *Diphyllobothrium*, *Strongyloides* in soil in Wasit province and the outcome of this survey is useful in conducting further studies in the future.

## References

**Afonso E, Lemoine M, Poulle M-L, Ravat M-C, Romand S, Thulliez P, Villena I, Aubert D, Rabilloud M, Riche B, Gilot- Fromont .** Spatial distribution of soil contamination by

**Anwar S , Naz A , Ashraf M Y , and Malik A.** Evaluation of inorganic contaminants emitted from automobiles and dynamics in soil, dust, and vegetations from major highways in Pakistan. *Environmental Science and Pollution Research*.2020; 27(26), 32494–32508.

**Badaki J A , Shitta K B , Labija G B., and Agwuja F S.** SOIL PARASITE CONTAMINATION OF PUBLIC PLACES WITHIN LOKOJA Badaki, J.A., Shitta, K Federal University Lokoja, Kogi State. Nigeria. *Bayero Journal of Pure and Applied Sciences* .2018; ISSN 2006 – 6996.

**Caldrer S , Ursini T, Santucci B , Motta L , and Angheben A. (2022).** Soil-Transmitted Helminths and Anaemia: A Neglected Association Outside the Tropics. *Microorganisms*.2022; 10(5), 1027.

**Collender PA, Kirby AE, Addiss DG, Freeman MC, Remais JV.** Methods for quantification of soil-transmitted helminthes in environmental media: Current techniques and recent advances. *Trends in Parasitology*.2015; 31, 625–639.

**Eisen AKA, Demoliner M, de Oliveira KG, Troian EA, Mallmann L, Filippi M, de Almeida PR, Spilki FR, Eisen AKA, Demoliner M, de Oliveira KG, Troian EA, Mallmann L, Filippi M, de Almeida PR, Spilki FR .** Soil contamination of a public park by human and canine mastadenovirus, as well as hookworms and *Toxocara* spp eggs. *Revista do Instituto de Medicina Tropical de São Paulo*.2019; 61, e60.

**Golek H I., and Al-Saeed A T M.** Contamination of soil with toxocara and other helminths in soils of Amadyia district, Duhok Governorate, Kurdistan Region-Iraq. *Applied Ecology & Environmental Research*2019; 17(6).

**Goshu A, Alemu G, Ayehu A .** Prevalence and intensity of soil transmitted helminths and associated factors among adolescents and adults in Bibugn Woreda, Northwest Ethiopia: a community based cross-sectional study. *J Trop Med* 2021:7043881.

**Hassan A A. Oyebamiji D A. and Idowu O F.** Spatial patterns of soil transmitted helminths in soil environment around Ibadan, an endemic area in South-west Nigeria. *Nigerian Journal of Parasitology*.2017; 38(2):179-184.

**Horiuchi S and Uga S .** Modified flotation method, an effective technique for recovering helminth eggs in soil. *Parasitol Int*. Oct;65(5 Pt B):576–579. doi: 10.1016/j.parint.2016.04.010. Epub 2016 May 1.

**Ihnacik L et al .** The survey of soil-transmitted helminth species abundance in Slovakia with an emphasis on parameters important for their distribution. *Front Med* 9.2022.

**Jourdan PM, Lamberton PHL, Fenwick A, Addiss DG.** Soil-transmitted helminth infections. *Lancet*.2018; 391(10117): 252–265.

**Kalkal H, Vohra S, Gupta S .** Anthelmintic resistance in livestock parasites: Indian scenario. *Int J Curr Microbiol*. 2020; App Sci 9(2):2839–2851

**KHAIRI S B , SHAKIR N N , and ABED M T.** A study of the parasitic contamination of commonly consumed vegetables in Wasit Province, Iraq. *Iranian Journal of Ichthyology* .2022; 9, 170-175.

**Koutsoumanis K, Allende A, Alvarez-Ordóñez A, Bolton D, Bover-Cid S, Chemaly M, Davies R, De Cesare A, Herman L, Hilbert F, Lindqvist R, NautaM, Peixe L, Ru G, Simmons M, Skandamis P, Suffredini E, Cacciò S, Chalmers R, Deplazes P, Devleesschauwer B, Innes E, Romig T, van der Giessen J, Hempen M, Van der Stede Y.** Robertson Public health risks associated with food-borne parasites. *EFSA Journal*.2018; 16, 5495.

**Loukas A , Maizels R M , and Hotez P J .** The yin and yang of human soil-transmitted helminth infections. *International Journal for Parasitology*.2021; 51(13–14).

**Masangay D U, Amado A J Y, Bulalas A R, Ciudadano S R , Fernandez J D , Sastrillo S M, and Mabaggu R M .** Association of Soil-transmitted Helminth Infection and Micronutrient Malnutrition: A Narrative Review. *Asian Journal of Biological and Life Sciences*.2021; 10(2), 317–324.

**Mohammed M A , Siddig E E , Elaagip A H , Edris M M A and Nasr A A.** Parasitic contamination of fresh vegetables sold at central markets in Khartoum Stat, Sudan. *Annals of Clinical Microbiology and Antimicrobials*.2016; 15:17 DOI 10.1186/s12941- 016- 0133-5.

**Montresor A et al .,** The global progress of soil-transmitted helminthiasis control in 2020 and World Health Organization targets for 2030. *PLoS Negl Trop Dis*.2020; 14(8):e0008505.

**Nath T C, Adnan M R , Sultana N , Husna A , Ndossi B A , Kang Y , Bia M M , Choe S , Park H, Lee D , Eamudomkarn C, Jeon H K , and Eom K S.** Integration of health education intervention to improve the compliance to mass drug administration for soil-transmitted helminths infection in Bangladesh: An implementation research. *Parasite Epidemiology and Control*. 2020; 11.

**Otero D, Alho AM, Nijse R, Roelfsema J, Overgaauw P, Madeira de Carvalho L .** Environmental contamination with *Toxocara* spp. eggs in public parks and playground sandpits of Greater Lisbon, Portugal. *Journal of Infection and Public Health*.2018; 11, 94–98.

**Otranto D, Deplazes P .** Zoonotic nematodes of wild carnivores. *International Journal for Parasitology: Parasites and Wildlife*.2019; 9, 370–383.

**Otranto D, Deplazes P.** Zoonotic nematodes of wild carnivores. *International Journal for Parasitology: Parasites and Wildlife*.2019; 9, 370–383.

**Prasetyo W A , and Ahmad A.** Iron-Deficiency Anemia Related to *Ancylostoma duodenale* Infection in 67 Years Old Female Patient. *Bioscientia Medicina : Journal of Biomedicine and Translational Research*. 2022; 6(3), 1455–1459.

**Raj E , Calvo-Urbano B , Heffernan C , Halder J , and Webster J P .** Systematic review to evaluate a potential association between helminth infection and physical stunting in children. *Parasites and Vectors*.2022; 15(1).

**Scott M E. and Koski K G .** Soil-Transmitted Helminths: Does Nutrition Make a Difference? In *Nutrition and Infectious Diseases* .2021; (pp. 325–364).

**Shapiro K, Bahia-Oliveira L, Dixon B, Dumètre A, de Wit LA, VanWormer E, Villena I.** Environmental transmission of *Toxoplasma gondii*: Oocysts in water, soil and food. *Food and Waterborne Parasitology*.2019; 15, e00049.

**Tadege B T , Mekonnen Z , Dana D , Sharew B , Dereje E , Loha E , Verweij J J, Casaert S , Vlamminck M A , Leveche B.** Assessment of environmental contamination with soil-transmitted helminths life stages at school compounds, households and open markets in Jimma Town, Ethiopia. *PLOS Neglected Tropical Diseases* .journal.pntd.0010307 April 4, 2022, pp 27.

**Thevenet PS, Alvarez HM, Torrecillas C, Jensen O, Basualdo JA..** Dispersion of *Echinococcus granulosus* eggs from infected dogs under natural conditions in Patagonia, Argentina. *Journal of Helminthology*2020; 94, e29, 1–7.

**Traversa D, Frangipane di Regalbono A, Di Cesare A, La Torre F, Drake J, Pietrobelli M.** Environmental contamination by canine geohelminths. *Parasites & Vectors*2014; 7, 67.

**Umhang G, Bastien M, Renault C, Faisse M, Caillot C, Boucher J-M, Hormaz V, Poulle M-L, Boué F. A.** flotation/sieving method to detect *Echinococcus multilocularis* and *Toxocara* spp. eggs in soil by real-time PCR. *Parasite*,2017.

**Utaaker KS, Skjerve E, Robertson LJ.** Keeping it cool: Survival of *Giardia* cysts and *Cryptosporidium* oocysts on lettuce leaves. *International Journal of Food Microbiology*.2017; 255,51–57.

**WHO (2023a)** Soil-transmitted helminth infections. [https:// www. who. int/ news- room/ fact- sheets/ detail/ soil- trans mitted- helmi nth- infections](https://www.who.int/news-room/fact-sheets/detail/soil-transmitted-helminth-infections). Accessed 17/11/2023.

**World Health Organization.** Reaching girls and women of reproductive age with deworming: Report of the WHO Advisory Group on deworming in girls and women of reproductive age. Geneva: World Health Organization; 2018. WHO/CDS/NTD/PCT/2018.01.

**Zeynudin A et al .,** Prevalence and intensity of soil-transmitted helminth infections and associated risk factors among household heads living in the peri-urban areas of Jimma town, Oromia, Ethiopia: a community-based cross-sectional study. *PLoS One*.2022; 17(9):e0274702.